

National Childhood and Adolescent Immunisation Schedule, Singapore For persons aged 0 to <18 years

Vaccination against	Birth	1 month	3 months	4 months	5 months	6 months	12 months	15 months	18 months	6-7 years ^A	10-11 years ^{AA}
Tuberculosis	BCG										
Hepatitis B*	HepB (D1)	HepB (D2)			HepB (D3) [#]						
Diphtheria, Tetanus, Pertussis			DTaP (D1)	DTaP (D2)	DTaP (D3)				DTaP (B1)		Tdap (B2)
Poliovirus			OPV (D1)	OPV (D2)	OPV (D3)				OPV (B1)	OPV (B2)	OPV (B3)
Measles, Mumps, Rubella							MMR (D1)	MMR (D2) ^{**}			
Pneumococcal Disease ^{**}			PCV (D1)		PCV (D2)		PCV (B1)				
Human Papillomavirus		<i>Recommended for females 9 to 26 years; three doses are required at intervals of 0, 2, 6 months</i>									
Influenza		<i>Recommended annually for all children aged 6 months to <5 years and children aged 6 months to <18 years in high-risk groups^{***}</i>									

Notes:

BCG Bacillus Calmette-Guérin
HepB Hepatitis B vaccine
DTaP Paediatric diphtheria and tetanus toxoid and acellular pertussis vaccine
Tdap Tetanus toxoid, reduced diphtheria toxoid and acellular pertussis vaccine
MMR Measles, mumps, and rubella vaccine
OPV Oral polio vaccine
PCV Pneumococcal conjugate vaccine
D1/D2/D3 1st dose, 2nd dose, 3rd dose
B1/B2/B3 1st booster, 2nd booster, 3rd booster
^A Primary 1
^{AA} Primary 5

3rd dose of HepB can be given with the 3rd dose of DTaP and OPV for the convenience of parents.
2nd dose of MMR can be given between 15-18 months
* Refer to 'Other Recommendations' below for guidelines on infants born to hepatitis B carrier mothers
** Refer to 'Other Recommendations' below for guidelines on catch-up vaccination schedule and vaccination for high-risk groups
*** Refer to 'Other Recommendations' below for more details on guidelines for influenza vaccination